Table of Contents

Title 48

PUBLIC HEALTH

GENERAL

Subpart 15.	Emergency Response Network	.1
Chapter 181.	General Provisions.	1
§18101.	Scope	1
Chapter 183.	Louisiana Emergency Response Network (LERN)	1
§18301.	Board Officers of Louisiana Emergency Response Network (LERN) Board	
§18303.	Quorum	
§18305.	Grounds for Removal of Board Members	
Chapter 185.	Regional Commissions; Membership; Officers; Meetings; Duties and Responsibilities	1
§18501.	Regional Commission Membership	
§18503.	Regional Commission Officers	
§18505.	Regional Commission Meetings	
§18507.	Regional Commission Duties and Responsibilities	2
Chapter 187.		
§18701.	Stroke Center Recognition	
§18703.	Stroke Center Criteria	3
§18705.	Attestation for Stroke Center Recognition	4
§18706.	Stroke Center Data Submission Requirements	4
§18708.	Failure to Submit Stroke Data to LERN	
Chapter 189.	Requirements for Louisiana STEMI Receiving/Referral Centers	5
§18901.	STEMI Center Recognition	
§18903.	Attestation for STEMI Center Recognition	5
§18905.	STEMI Center Listing	
§18907.	Hospital Destination/STEMI System Transport:	5
Chapter 191.		
§19101.	Entry Criteria and Region 4 LERN LCC Destination Protocol	5
§19103.	Region 7 LERN Entry and Destination Protocols	6
§19105.	Standard LERN Entry Criteria; Standard Destination Protocol	8
§19107.	Interregional Transfer Protocol	9
§19109.	Standard LERN Entry and Destination Criteria	10
§19111.	Interregional Transfer Protocol	11
§19113.	LERN Entry Criteria: Trauma; LERN Destination Protocol: Trauma	12
§19115.	LERN Destination Protocol: TRAUMA	13
§19117.	LERN Destination Protocol: Trauma	13
§19119.	Destination Protocol: TRAUMA	14
§19121.	LERN Destination Protocol: TRAUMA	15
Chapter 192.	Burn Protocols	16
§19201.	LERN Destination Protocol: BURN	16
Chapter 193.	Stroke Protocols	16
§19301.	LERN Destination Protocol: Stroke	16
§19303.	LERN Destination Protocol: Stroke	17

Chapter 195.	STEMI Protocols	17
§19501.	STEMI Triage Protocol for Pre-Hospital Providers	17
Chapter 197.	Trauma Program Recognition	18
_	Generally	
	Purpose	
§19705.	Qualifications for LERN Trauma Program Recognition	18
§19707.	Procedure for Trauma Program Recognition	

This page left intentionally blank

Subpart 15. Emergency Response Network

Chapter 181. General Provisions

§18101. Scope

A. These rules are adopted by the Louisiana Emergency Network (hereinafter LERN) Board (hereinafter board) to effectuate the provisions of R.S. 40:2841 et seq.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Louisiana Emergency Response Network Board, LR 34:649 (April 2008).

Chapter 183. Louisiana Emergency Response Network (LERN)

§18301. Board Officers of Louisiana Emergency Response Network (LERN) Board

- A. The chairman and vice-chairman, and any other officers that the board shall deem necessary, shall be elected for a two-year term at the first meeting held following January 1 of each even numbered year.
- B. In the case of a vacancy in the office of chairman, the vice-chairman shall serve the remainder of the vacated term, and in the case of a vacancy in the office of vice-chairman, the board shall elect a new vice-chairman who shall serve the remainder of the vacated term.

C. The chairman shall:

- 1. preside at all meetings of the board;
- 2. determine necessary subcommittees and working group and appoint members to each subcommittee and working groups;
 - 3. direct activities of staff between board meetings;
- 4. provide direction on behalf of board between meetings to all regional commissions;
- 5. designate the date, time and place of board meetings;
- 6. enter into confidentiality agreements on behalf of the board regarding pertinent data to be submitted to board and board staff which contain individually identifiable health or proprietary information;
- 7. perform all other duties as may be assigned by the board.
- D. Should the chairman become unable to perform the duties of chairman, the vice-chairman shall act in his stead.
- E. A ground for removal of a board officer includes conviction of a felony.

AUTHORITY NOTE: Promulgated in accordance with R.S. 48:2844(H) and 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Louisiana Emergency Response Network Board, LR 34:650 (April 2008).

§18303. Quorum

A. Eight members of the board shall constitute a quorum for all purposes.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2844(H) and 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Louisiana Emergency Response Network Board, LR 34:650 (April 2008).

§18305. Grounds for Removal of Board Members

A. Grounds for removal of board members include conviction of a felony.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2844(H) and 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Louisiana Emergency Response Network Board, LR 34:650 (April 2008).

Chapter 185. Regional Commissions; Membership; Officers; Meetings; Duties and Responsibilities

§18501. Regional Commission Membership

- A. Selection of Regional Commission Membership by Louisiana Emergency Response Network (LERN) Board
- 1. The process for selecting the regional commission members is as follows:
- a. the LERN Board Chairman shall request in writing the name of a nominee to serve on each regional commission from each of the legislatively identified state organizations;
- b. in the event there is more than one organization, state association or entity, each entity shall be requested to name a nominee and, once constituted, the commission shall choose from among the nominees; and
- c. if no state or local organization exists in a category, but multiple nominees are identified in that category, the selection of the representative to serve on the regional commission will be determined by that category's group of nominees.
- 2. Once documentation is received from each organization or group, the compiled list of nominees is submitted to the board for ratification. The board shall appoint those selected by the various organizations.
- B. Voting members of the regional commission may be added through a process employing the following steps:
- 1. majority vote of a quorum of voting members of the commission;
- 2. formal written request to LERN Board to add specified voting member, with reasons for adding. Such addition must represent a group which would enhance the working of the regional commission;

- 3. majority vote by LERN Board members at a meeting. If such a vote fails, the regional commission may appear in person at the following LERN Board meeting, where the subject will be revisited;
- 4. once an additional voting member is approved for one region, in order for other regions to add a member representing the same group, only a letter detailing the requirements of Paragraphs 1 through 3 above will be necessary to add the particular member. Board approval will not require an additional vote.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2845(A)(3)(a) and 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Louisiana Emergency Response Network Board, LR 34:650 (April 2008).

§18503. Regional Commission Officers

- A. Each regional commission shall select a chairman and vice chairman.
- B. The chairman and vice-chairman, and any other officers that the commission shall deem necessary, shall be elected for a two-year term at the first meeting held following January 1 of each even numbered year.
- C. In the case of a vacancy in the office of chairman, the vice-chairman shall serve as chairman for the remaining vacated term; and in the case of a vacancy in the office of vice-chairman, the regional commission shall elect a new vice-chairman who shall serve until the expiration of the vacated term.
 - D. The chairman shall:
 - 1. preside at all meetings of the commission;
- 2. determine necessary ad hoc committees, appoint a commission member to chair each such committee, and provide for the commission as a whole to name the membership of the committee;
- 3. provide direction to the commission to implement the mandates of the LERN Board;
- 4. direct that a record of all meetings of the commission shall be kept and such records shall be retained as permanent records of the transactions of the commission; and
- 5. perform all other duties pertaining to the office of chairman of the commission or as may be assigned by the commission.
- E. Should the chairman become unable to perform the duties of chairman, the vice-chairman shall act in his stead.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2845(A)(3)(a) and 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Louisiana Emergency Response Network Board, LR 34:650 (April 2008).

§18505. Regional Commission Meetings

A. Meetings of the commission shall be noticed, convened and held not less frequently than quarterly during

each calendar year and otherwise at the call of the chairman or on the written petition for a meeting signed by not less than the number of members which would constitute a quorum of the commission. Meetings shall be held on such date and at such time and place as may be designated by the chairman.

B. One third of the currently serving members of the commission shall constitute a quorum for all purposes. All actions which the commission is empowered by law to take shall be effected by vote of not less than a majority of the members present at a meeting of the commission at which a quorum is present.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2845(A)(3)(a) and 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Louisiana Emergency Response Network Board, LR 34:651 (April 2008).

§18507. Regional Commission Duties and Responsibilities

- A. Each regional commission shall:
- 1. develop a written system plan for submission to LERN Board, which plan shall:
- a. identify all resources available in the region for emergency and disaster preparedness and response;
- b. be based on standard guidelines for comprehensive system development;
- c. include all parishes within the region unless a specific parish portion thereof has been aligned within an adjacent region;
- d. give an opportunity to all health care entities and interested specialty centers opportunity to participate in the planning process; and
 - e. address the following components:
 - i. injury prevention;
 - ii. access to the system;
 - iii. communications;
 - iv. pre-hospital triage criteria;
 - v. diversion policies;
 - vi. bypass protocols;
 - vii. regional medical control;
 - viii. facility triage criteria;
 - ix. inter-hospital transfers;
- x. planning for the designation of trauma facilities, including the identification of the lead facility(ies); and
- xi. a performance improvement program that evaluates processes and outcomes from a system perspective;

- 2. upon approval of the board, implement the system plan to include:
- a. education of all entities about the plan components;
- b. on-going review of resource, process, and outcome data; and
- c. if necessary, revision and re-approval of the plan or plan components by LERN Board;
- 3. annually complete a regional needs assessment and conduct education and training within the region to meet the needs identified in the annual needs assessment;
- 4. develop and implement a regional performance improvement (PI) program plan;
- 5. develop and implement a regional injury prevention program;
- 6. at least quarterly, submit evidence of on-going activity, including meeting notices and minutes, to LERN Board; and
- 7. Annually submit a report to LERN Board which describes progress toward system development and demonstrates on-going activity;
- B. Regional commission may request technical assistance from the LERN Board at any time.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2845(A)(3)(a) and 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Louisiana Emergency Response Network Board, LR 34:651 (April 2008).

Chapter 187. Requirements for Louisiana Stroke Center Recognition

§18701. Stroke Center Recognition

- A. The Louisiana Emergency Response Network Board (LERN) and the Louisiana Department of Health recognize the following six levels of stroke facilities:
- 1. CSC: comprehensive stroke center (formerly designated as level 1);
 - 2. TSC: thrombectomy capable stroke center;
- 3. PSC-E: primary stroke center with endovascular capability;
- 4. PSC: primary stroke center (formerly designated as level 2);
- 5. ASRH: acute stroke ready hospital (formerly designated as level 3); and
- 6. stroke bypass hospital (formerly designated as level 4).
- B. Participation in Louisiana stroke center recognition is voluntary and no hospital shall be required to participate.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2846(A) and 48:2845(A)(7).

HISTORICAL NOTE: Promulgated by the Department of Health, Emergency Response Network, LR 40:2590 (December 2014), amended by the Department of Health, Emergency Response Network LR 46:1088 (August 2020).

§18703. Stroke Center Criteria

- A. Each facility participating in stroke center recognition shall meet the following criteria.
- 1. CSC: a comprehensive stroke center (CSC) will meet the requirements specified by the joint commission or other board approved accrediting/certification body approved by LERN for comprehensive stroke center certification. Attestation as a CSC is only allowed after verification by the joint commission or other LERN approved accrediting/certification body that the facility meets all requirements set forth in the CSC standards.
- 2. TSC: a thrombectomy capable stroke center (TSC) will meet the requirements specified by the joint commission or other board approved accrediting/certification body approved by LERN for thrombectomy capable stroke center certification. Attestation as a TSC is only allowed after verification by the joint commission or other LERN approved accrediting/certification body that the facility meets all requirements set forth in the TSC standards.
- 3. PSC-E: a primary stroke center (PSC-E) shall meet the requirements specified by the joint commission, healthcare facilities accreditation program (HFAP), or other LERN approved accrediting/certification body for Primary Stroke Center verification. Attestation as a PSC-E is only allowed after verification by the joint commission, HFAP, or other LERN approved accrediting/certification body that the facility meets all requirements set forth in the PSC standards. In addition to PSC requirements, a PSC-E must have physician(s) credentialed to perform mechanical thrombectomy and must update resource management portal of endovascular availability at all times. If a physician credentialed to perform endovascular capability is not available, the PSC-E must notify all EMS providers in the region when endovascular resources are not available. The PSC-E must collect and submit quarterly to LERN the same data the joint commission requires the Thrombectomy Stroke Capable centers to collect.
- 4. PSC: a primary stroke center (PSC) shall meet the requirements specified by the joint commission, healthcare facilities accreditation program (HFAP), or other LERN approved accrediting/certification body for primary stroke center verification. Attestation as a PSC is only allowed after verification by the joint commission, HFAP, or other LERN approved accrediting/certification body that the facility meets all requirements set forth in the PSC standards.
- 5. ASRH: an acute stroke ready hospital (ASRH) will provide timely access to stroke care but may not meet all criteria for a CSC, TSC, or a PSC or a PSC-E facility. An ASRH will provide acute stroke care in urban and rural areas where transportation and access are limited. An ASRH is intended to recognize models of care delivery that have shown utility, including "drip-and-ship" and telemedicine. An ASRH must meet requirements adopted by LERN and

submit quarterly data as required by LERN. LERN approved requirements are based on national best practice guidelines.

- 6. Stroke bypass hospital: a stroke bypass hospital should not receive patients exhibiting signs or symptoms of stroke except for instances when the clinical situation requires stopping at the closest emergency department. A stroke bypass hospital must have:
- a. transfer protocol in place for transfer to higher levels of care through written and agreed upon relationship with a CSC, TSC, PSC, PSC-E or ASRH stroke center.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2846(A) and 48:2845(A)(7).

HISTORICAL NOTE: Promulgated by the Department of Health, Emergency Response Network, LR 40:2590 (December 2014), amended by the Department of Health, Emergency Response Network LR 46:1088 (August 2020).

§18705. Attestation for Stroke Center Recognition

- A. A hospital seeking CSC, TSC, PSC-E, ASRH or stroke bypass recognition will submit an affidavit of the hospital CEO to LERN detailing compliance with the requirements designated herein.
- 1. A center or hospital seeking CSC recognition which submits a copy of that level of certification by a LERN-recognized organization, such as the joint commission or other LERN approved accrediting/certification body, shall be assumed to meet the requirements for recognition.
- 2. A center or hospital seeking TSC stroke center recognition which submits a copy of that level of certification by a LERN-recognized organization, such as the joint commission, HFAP, or other LERN approved accrediting/certification body, shall be assumed to meet the requirements for recognition.
- 3. A center or hospital seeking PSC-E stroke center recognition which submits a copy of PSC certification by a LERN-recognized organization, such the joint as commission, HFAP, other **LERN** approved or accrediting/certification body, shall be assumed to meet the requirements for recognition. In addition to a copy of the certification, the CEO must also attest to meeting the additional board approved requirements.
- 4. A center or hospital seeking PSC stroke center recognition which submits a copy of that level of certification by a LERN-recognized organization, such as the joint commission, HFAP, or other LERN approved accrediting/certification body, shall be assumed to meet the requirements for recognition.
- 5. Although a center or hospital seeking ASRH stroke center recognition is not required to obtain certification by an external certifying body, a hospital which submits a copy of ASRH certification by a LERN-recognized organization, such as the joint commission, HFAP or other LERN approved accrediting/certification body, shall be assumed to meet the requirements for recognition. Hospitals must all meet LERN ASRH requirements and approved data submission requirements.

- 6. Each center or hospital shall submit proof of continued compliance every two years by submission of an affidavit by its CEO.
- B. A hospital or center which fails to meet the requirements as attested, or which no longer chooses to maintain state Stroke Facility level recognition, shall immediately notify LERN and local EMS.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2846(A) and 48:2845(A)(7).

HISTORICAL NOTE: Promulgated by the Department of Health, Emergency Response Network, LR 40:2590 (December 2014), amended by the Department of Health, Emergency Response Network LR 46:1089 (August 2020).

§18706. Stroke Center Data Submission Requirements

- A. All stroke centers, whether CSC, TSC, PSC-E, PSC or ASRH are required to submit certain data to the board on a quarterly basis.
- B. The requirements of and for data submission are posted on the LERN website, http://lern.la.gov/lern-stroke-system/stroke-data-collection.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2846(A) and 48:2845(A)(7).

HISTORICAL NOTE: Promulgated by the Department of Health, Emergency Response Network, LR 46:1089 (August 2020).

§18708. Failure to Submit Stroke Data to LERN

- A. Acute stroke ready hospitals not submitting data for one quarter or not submitting the required action plan and/or mock code, if applicable, will result in automatic probation, which will generate a warning letter to the CEO. The letter will communicate LERN board expectation for data and (action plan and/or mock code, if applicable) submission for the missed quarter and the following quarter.
- B. For an ASRH not submitting data to the board for two consecutive quarters, the hospital will automatically be demoted to a stroke bypass hospital.
- C. Once an ASRH demotes to a stroke bypass hospital for non-adherence with submission requirement, the hospital CEO cannot re-attest until the hospital has submitted two consecutive quarters of data meeting standards.
- D. If an ASRH fails to meet the performance metrics after two quarters of participation in data review, the board appointed stroke subcommittee will present the blinded data to the board for a vote on demotion to stroke bypass hospital versus continued remediation.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2846(A) and 48:2845(A)(7).

HISTORICAL NOTE: Promulgated by the Department of Health, Emergency Response Network, LR 46:1089 (August 2020).

Chapter 189. Requirements for Louisiana STEMI Receiving/Referral Centers

§18901. STEMI Center Recognition

- A. The Louisiana Emergency Response Network Board (LERN), and the Louisiana Department of Health and Hospitals recognize the following types of facilities for the treatment of ST elevated myocardial infarction (STEMI):
 - 1. STEMI receiving center; and
 - 2. STEMI referral center.
- B. Participation in the Louisiana STEMI center recognition is voluntary and no hospital shall be required to participate.
- C. A facility seeking STEMI receiving center recognition shall meet the STEMI receiving center requirements adopted by LERN. LERN approved requirements are based on national best practice guidelines.
- D. A hospital with an emergency room not meeting criteria for a STEMI receiving center will automatically default to a STEMI referral center.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2846(A) and 48:2845(A)(7).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 40:2591 (December 2014).

§18903. Attestation for STEMI Center Recognition

- A. A hospital seeking STEMI Center recognition will submit an affidavit of the hospital CEO to LERN detailing compliance with LERN Approved STEMI Receiving center requirements.
- 1. Those hospitals which submit a copy of certification by a LERN-recognized organization such as The American Heart Association Mission:Lifeline, Society of Cardiovascular Patient Care or other LERN approved accrediting/certification body shall be assumed to meet the requirements for recognition.
- 2. Each center or hospital shall submit proof of continued compliance every two years by submission of an affidavit of its CEO.
- B. A hospital or center which fails to meet the criteria for a STEMI receiving center or which no longer choose to maintain state STEMI receiving center recognition, shall immediately notify LERN and local EMS.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2846(A), 48:2845(A)(7) and R.S. 9:2798.5.

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 40:2591 (December 2014).

§18905. STEMI Center Listing

A. LERN will publish a list on its website of hospitals or centers attesting to STEMI center criteria for recognition as either a STEMI receiving center or STEMI referral center.

This list shall be made available to the LERN regional commissions for facilitation of EMS transportation plans.

AUTHORITY NOTE: Promulgated in accordance with La. R.S. 40:2846(A) and 48:2845(A)(7).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 40:2591 (December 2014).

§18907. Hospital Destination/STEMI System Transport:

- A. These rules are not intended to prevent any hospital or medical facility from providing medical care to any patient but rather to serve as a guideline to facilitate the timely and appropriate delivery of STEMI patients to the most appropriate care site for the definitive treatment of STEMI.
- B. Knowledge of STEMI capabilities and the use of a STEMI pre-hospital destination protocol will enable providers to make timely decisions, promote appropriate utilization of the STEMI care delivery system, and ultimately save lives.

AUTHORITY NOTE: Promulgated in accordance with La. R.S. 40:2846(A), 48:2845(A)(7) and R.S. 9:2798.5.

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 40:2591 (December 2014).

Chapter 191. Trauma Protocols

§19101. Entry Criteria and Region 4 LERN LCC Destination Protocol

A. On November 15, 2007, the Louisiana Emergency Response Network Board [R.S. 40:2842(1)] adopted and promulgated "LERN Entry Criteria" and "LERN Region 4 LCC Destination Protocol" for region 4 of the Louisiana Emergency Response Network (R.S. 40:2842(3)), which region includes the parishes of Acadia, Evangeline, Iberia, Lafayette, St. Martin, St. Landry, and Vermilion, as follows.

1. LERN Entry Criteria

LERN Entry Criteria			
□ Unmanageable Airway □ Tension Pneumothorax □ Traumatic cardiac arrest □ Burn patient without patent airway □ Burn patient >40% BSA without IV	YES→	Call LCC	
Neurologic Trauma GCS <14 + one or more of the following: Penetrating head injury or depressed skull fracture Open head injury with or without CSF leak Deterioration of the GCS Lateralizing signs or paralysis (i.e., one-sided weakness, motor, or sensory deficit)	YES→	Call LCC	
Physiologic □ SBP <90 (adults and > 9 y/o) <70 + 2 [age (yrs)] (age 1 to 8) <70 (age 1 to 12 months) <60 (term neonate) □ RR <10 or >29 (adults and > 9 y/o) <15 or >30 (age 1 to 8) <25 or >50 (<12 m/o) Anatomic	YES→	Call LCC	

	LERN Entry Criteria				
	All penetrating injuries to neck, torso and				
	extremities proximal to elbow and knee				
	Flail Chest				
	2 or more proximal long-bone fractures				
	Crush, degloved or mangled extremity				
	Amputation proximal to wrist and ankle				
	Pelvic Fracture				
	Hip fractures (hip tenderness, deformity, lateral deviation of foot)	$YES{\rightarrow}$	Call LCC		
	Major joint dislocations (hip, knee, ankle, elbow)				
lп	Open Fractures				
lä	Fractures with neurovascular compromise				
	(decreased peripheral pulses or prolonged				
	capillary refill, motor or sensory deficits				
	distal to fracture, etc.)				
Me	echanism				
	Falls > 20 ft. (adults)				
	> 10 ft. (child) or 2 to 3 times height				
	High-risk auto crash				
	Intrusion > 12 in. occupant site:				
	>20 in. any site		Call		
	Ejection, partial or complete from automobile	YES→	LCC		
	Death in same passenger compartment				
	Auto vs. pedestrian/bicyclist thrown, run				
	over or >5 MPH impact				
	Motorcycle crash >20 MPH				
Spe	ecial				
	Pregnancy >20 weeks	YES→	Call		
	Burns (will follow ABA guidelines)	1 Lo→	LCC		
Otl	Other				
	Age \geq 55 y/o or \leq 8 y/o				
	Anticoagulation and bleeding disorders	YES→	Call		
	End stage renal disease	1 Lo→	LCC		
	Transplant patients				

2. LERN Region 4 LCC Destination Protocol

LERN Region 4 LCC Destination Protocol			
 □ Unmanageable Airway □ Tension Pneumothorax □ Traumatic cardiac arrest □ Burn patient without patent airway □ Burn patient >40 percent BSA without IV 	YES→	Closest ED	
NO			
Neurologic Trauma			
□ GCS <14 + one or more of the following: □ Penetrating head injury or depressed skull fracture □ Open head injury with or without CSF leak □ Deterioration of the GCS □ Lateralizing signs or paralysis (i.e., one-sided weakness, motor, or sensory deficit)	YES→	LERN Level II	
NO ↓			
Physiologic			
SBP <90 (adults and > 9 y/o) <70 + 2 [age (yrs)] (age 1 to 8) <70 (age 1 to 12 months) <60 (term neonate) RR <10 or >29 (adults and ≥ 9 y/o) <15 or >30 (age 1 to 8) <25 or >50 (<12 m/o)	YES→	LERN Level II or III	
NO ↓	•		

	LERN Region 4 LCC Destination Protocol				
An	atomic				
	All penetrating injuries to neck, torso and extremities proximal to elbow and knee Flail Chest 2 or more proximal long-bone fractures Crush, degloved or mangled extremity Amputation proximal to wrist and ankle Pelvic Fracture Hip fractures (hip tenderness, deformity, lateral deviation of foot) Major joint dislocations (hip, knee, ankle, elbow) Open Fractures Fractures with neurovascular compromise (decreased peripheral pulses or prolonged capillary refill, motor or sensory deficits	YES→	LERN Level II or III		
	distal to fracture, etc.)				
	↓				
Me	echanism	,			
	Falls > 20 ft. (adults) > 10 ft. (child) or 2 to 3 times height High-risk auto crash Intrusion > 12 in. occupant site: > 20 in. any site Ejection, partial or complete from automobile Death in same passenger compartment Auto vs. pedestrian/bicyclist thrown, run over or > 5 MPH impact Motorcycle crash > 20 MPH	YES→	LERN Level II or III		
	NO J				
Sn	ecial				
	Pregnancy >20 weeks Burns (will follow ABA guidelines)	YES→	LERN		
	NO ↓				
Ot	Other				
	Age ≥ 55 y/o or <8 y/o Anticoagulation and bleeding disorders End stage renal disease Transplant patients	YES→	LERN Level II, III or IV		

B. On June 26, 2008, the Louisiana Emergency Response Network Board passed a resolution allowing any region of the Louisiana Emergency Response Network which agreed to use the foregoing "LERN Entry Criteria" and "LERN Region 4 LCC Destination Protocol" to begin operating using the "LERN Entry Criteria" and "LERN Region 4 LCC Destination Protocol" set forth above.

C. This protocol was published at LR 35:1181-1183 (June 20, 2009).

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 41:138 (January 2015).

§19103. Region 7 LERN Entry and Destination Protocols

A. On November 15, 2007, the Louisiana Emergency Response Network Board [R.S. 40:2842(1)] adopted and promulgated "Region 7 LERN Entry and Destination Protocol" for region 7 of the Louisiana Emergency Response Network [R.S. 40:2842(3)], which region includes the

parishes of Bienville, Bossier, Caddo, Claiborne, DeSoto, Natchitoches, Red River, Sabine and Webster, as follows.

- 1.a. Traumatic patients who meet the following criteria will be entered to LERN call center and should be transported directly to LSUHSC in Shreveport, Louisiana, if possible:
- i. airway compromise (intubated, apneic, or obstructed airway);
- ii. penetrating wound of head, neck, chest, abdomen, groin, or buttocks;
 - iii. blood pressure ≤ 100 or signs of shock;
 - iv. GCS 12 or less;
- v. new onset neurological deficit associated with traumatic event;
- vi. extremity wound with absent pulse or amputation proximal to foot or hand.
- b. Trauma patients who meet the following criteria, and are located outside the city limits of Shreveport and Bossier City, should be taken to nearest hospital for immediate stabilization followed by continued rapid transport to LSUHSC Shreveport per the LERN hospital protocol:
- i. unable to establish and maintain adequate airway/ventilation;
- ii. hypotension unresponsive to crystalloids (no more than $2\ L$);
- iii. patients who meet trauma center criteria but have a transport time > 60 minutes;
 - iv. traumatic arrest.
- B. On May 8, 2008, the Louisiana Emergency Response Network Board (R.S. 40:2842(1)) amended and promulgated, as amended, "Region 7 LERN Entry and Destination Protocol" for region 7 of the Louisiana Emergency Response Network (R.S. 40:2842(3)), which region includes the parishes of Bienville, Bossier, Caddo, Claiborne, DeSoto, Natchitoches, Red River, Sabine and Webster, which protocol was originally adopted and promulgated on November 15, 2007, so that the "Region 7 Louisiana Emergency Response Network Entry and Destination Protocol," as amended, effective May 8, 2008, is as follows.
- 1.a. Traumatic patients who meet the following criteria will be entered to LERN call center and should be transported directly to LSUHSC in Shreveport, if possible:
- i. airway compromise (intubated, apneic, or obstructed airway);
- ii. penetrating wound of head, neck, chest, abdomen, groin, or buttocks;
 - iii. blood pressure ≤ 100 or signs of shock;
 - iv. GCS 12 or less;

- v. new onset neurological deficit associated with traumatic event;
- vi. extremity wound with absent pulse or amputation proximal to foot or hand;
- vii. burn patients as identified following ABA guidelines;
- viii. healthcare provider discretion—patients evaluated by hospitals may be entered into LERN if the evaluating hospitals medical personnel determines the patient has a medical condition requiring immediate surgical evaluation and/or intervention and the transferring hospital does not have these services immediately available at that facility (Healthcare provider discretion does not include orthopedic injuries.).
- b. Patients that have been entered into LERN but will require greater than 60 minute transport time from the field should stop at local area hospitals for stabilization. These patients should still be entered into LERN from the field but will require transport to local area hospitals for stabilization. LERN will facilitate the movement of these patients from the local hospital once stabilizing measures are completed.
- i. The following are conditions requiring immediate stabilization by local area hospitals:
- (a). unable to establish and maintain adequate airway/ventilation;
- (b). hypotension unresponsive to crystalloids (no more than $2\ L$);
- (c). patients who meet trauma center criteria but have a transport time > 60 minutes;
 - (d). traumatic arrest.
- C. The following will be routed directly to the LSUHSC Burn Unit from local area hospitals or from the field:
- 1. partial-thickness and full thickness burns greater than 10 percent of the total body surface area (TBSA) in patients younger than 10 years of age or older than 50 years of age;
- 2. partial-thickness and full thickness burns greater than 20 percent of the total body surface area (TBSA) in other age groups;
- 3. partial-thickness and full thickness burns involving the face, eyes, ears, hands, feet, genitalia, perineum, or skin overlying major joints;
- 4. full-thickness burns greater than 5 percent TBSA in any age group;
 - 5. electrical burns, including lightning injury;
 - 6. chemical burns;
 - 7. patients with inhalation injury;

- 8. burn injury in patients with pre-existing illnesses that could complicate management, prolong recovery, or adversely affect mortality risk;
- 9. any burn patient in whom concomitant trauma poses an increased risk of morbidity or mortality may be treated initially in a trauma center until stable before transfer to a burn center:
- 10. children with burns seen in hospitals without qualified personnel or equipment for their care;
- 11. burn injury in patients who will require special social and emotional or long-term rehabilitative support, including cases involving suspected child abuse or neglect.
- D. These protocols were published at LR 35:1183-1184 (June 20, 2009).

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 41:139 (January 2015).

§19105. Standard LERN Entry Criteria; Standard Destination Protocol

A. On June 18, 2009, the Louisiana Emergency Response Network Board (R.S. 40:2842(1)) adopted and promulgated "Standard LERN Entry Criteria" and "Standard Destination Protocol" for use in all regions, except region 7, of the Louisiana Emergency Response Network (R.S. 40:2842(3)), replacing the "LERN Entry Criteria" and "LERN Region 4 LCC Destination Protocol," adopted and promulgated November 15, 2007, as follows.

1. Standard LERN Entry Criteria—Pre-Hospital and Hospital Triage Protocol

Standard LERN Entry Criteria				
Pre-Hospital and Hospital Triage Pr	otocol			
☐ Unmanageable Airway				
☐ Tension Pneumothorax		Call		
☐ Traumatic cardiac arrest	$YES \rightarrow$	LCC		
☐ Burn patient without patent airway		LCC		
☐ Burn patient >40 percent BSA without IV				
Neurologic Trauma				
\Box GCS <14 + one or more of the following:				
 Penetrating head injury or depressed skull 				
fracture		Call		
☐ Open head injury with or without CSF leak	$YES \rightarrow$	LCC		
☐ Deterioration of the GCS		LCC		
☐ Lateralizing signs or paralysis (i.e., one-sided				
weakness, motor, or sensory deficit)				
Physiologic				
\square SBP <90 (adults and > 9 y/o)				
<70 + 2 [age (yrs)] (age 1 to 8)				
<70 (age 1 to 12 months)		Call		
<60 (term neonate)	$YES \rightarrow$	LCC		
\square RR <10 or >29 (adults and > 9 y/o)		LCC		
<15 or >30 (age 1 to 8)				
<25 or >50 (<12 m/o)				
Anatomic				
☐ All penetrating injuries to neck, torso and				
extremities proximal to elbow and knee		Call		
☐ Flail Chest	$YES \rightarrow$	LCC		
☐ 2 or more proximal long-bone fractures		LCC		
☐ Crush, degloved or mangled extremity				

Standard LERN Entry Criteria				
	Pre-Hospital and Hospital Triage Protocol			
	Amputation proximal to wrist and ankle			
	Pelvic Fracture			
	Hip fractures (hip tenderness, deformity, lateral			
	deviation of foot)			
	Major joint dislocations (hip, knee, ankle,			
	elbow)			
	Open Fractures			
	Fractures with neurovascular compromise			
i	(decreased peripheral pulses or prolonged			
	capillary refill, motor or sensory deficits distal			
	to fracture, etc.)			
Me	echanism			
Ш	Falls > 20 ft. (adults)			
	> 10 ft. (child) or 2 to 3 times height			
	High-risk auto crash Intrusion > 12 in, occupant site:			
	>18 in. any site			
i	Ejection, partial or complete from		Call	
	automobile	$YES \rightarrow$	LCC	
	 Death in same passenger 		LCC	
	compartment			
П	Auto vs. pedestrian/bicyclist thrown, run over or			
	significant (>20 MPH) impact			
	Motorcycle crash >20 MPH			
Sp	ecial			
	Pregnancy >20 weeks	MEG .	Call	
	Burns (will follow ABA guidelines)	$YES \rightarrow$	LCC	
Ot	her			
	Age ≥ 55 y/o or <8 y/o			
	Anticoagulation and bleeding disorders	YES→	Call	
	End stage renal disease	1 E3→	LCC	
	Transplant patients			

2. Standard Destination Protocol

Standard Destination Protocol				
☐ Unmanageable Airway				
☐ Tension Pneumothorax		Closest		
☐ Traumatic cardiac arrest	$YES \rightarrow$	ED		
☐ Burn patient without patent airway		ED		
☐ Burn patient >40 percent BSA without IV				
NO				
\downarrow				
Neurologic Trauma				
\Box GCS <14				
☐ Penetrating head injury or depressed skull				
fracture		LERN		
☐ Open head injury with or without CSF leak	YES→	Level I		
☐ Deterioration of the GCS	1L5-7	or II		
☐ Lateralizing signs or paralysis		01 11		
(i.e., one-sided weakness, motor, or sensory				
deficit)				
NO				
↓				
Physiologic				
\Box SBP <90 (adults and >9 y/o)				
<70 + 2 [age (yrs)] (age 1 to 8)				
<70 (age 1 to 12 months)		LERN		
<60 (term neonate)	$YES \rightarrow$	Level I,		
\square RR <10 or >29 (adults and >9 y/o)		II or III		
<15 or >30 (age 1 to 8)				
<25 or >50 (<12 m/o)				
NO				
↓				
Anatomic				

	Standard Destination Protocol			
	All penetrating injuries to neck, torso and			
	extremities proximal to elbow and knee			
	Flail Chest			
	2 or more proximal long-bone fractures			
	Crush, degloved or mangled extremity			
	Amputation proximal to wrist and ankle			
	Pelvic Fracture		LERN	
	Hip fractures (hip tenderness, deformity, lateral	$YES \rightarrow$	Level I,	
	deviation of foot)		II or III	
	Major joint dislocations (hip, knee, ankle, elbow)			
	Open Fractures			
	Fractures with neurovascular compromise			
	(decreased peripheral pulses or prolonged			
	capillary refill, motor or sensory deficits distal			
	to fracture, etc.)			
	NO			
	↓			
Me	echanism			
	Falls >20 ft. (adults)			
	>10 ft. (child) or 2 to 3 times height			
	High-risk auto crash			
	 Intrusion >12 in. occupant site: 			
	• >18 in. any site		LERN	
	 Ejection, partial or complete from 	$YES \rightarrow$	Level	
	automobile		II or III	
	 Death in same passenger compartment 			
	Auto vs. pedestrian/bicyclist thrown, run over or			
	significant (>20 MPH) impact			
	Motorcycle crash >20 MPH			
	NO			
	\downarrow			
Sp	ecial			
	Pregnancy >20 weeks		LERN	
	Burns (will follow ABA guidelines)	YES→	Level II or III	
	NO			
	\downarrow			
Other				
	$Age \ge 55 \text{ y/o or } < 8 \text{ y/o}$		LERN	
	Anticoagulation and bleeding disorders	MEG	Level	
	End stage renal disease	$YES \rightarrow$	II, III	
	Transplant patients		or IV	

B. These protocols were published at LR 35:1409 (July 20, 2009).

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 41:140 (January 2015).

§19107. Interregional Transfer Protocol

- A. On June 18, 2009, the Louisiana Emergency Response Network Board [R.S. 40:2842(1)] adopted and promulgated "Interregional Transfer Protocol" for the Louisiana Emergency Response Network [R.S. 40:2842(3)], as follows.
- 1. The LERN interregional transfer protocol only applies to those regions and (hospitals/EMS) that are participating in the LERN network.
- 2. The interregional transfer protocol will be tested over a 90 day period. At the end of the 90 days all interregional transfers will be reviewed for compliance with

protocols, quality, patient safety and standards of care. This information will be shared with commissions of the regions participating as well as the LERN board and the "design the system group". Decisions regarding the interregional transfer protocol will be made at the end of the 90 days trial period.

3. Interregional Transfer Protocol

- a. All patients whose condition exceeds the regionally available resources provided by local area hospitals may be transferred from one region to another following LERN interregional transfer protocol. Destination to the definitive care hospital in the receiving region will follow the LERN standard protocol (all laws regarding EMTALA apply).
- b. Only regions operating with the LERN standard protocol will be involved in the LERN interregional transfer protocol.
- c. Patients being transferred via the LERN interregional transfer protocol must:
- i. be assessed at a local area hospital for treatment and stabilized by a physician and meet the entry criteria as determined by LERN standard protocol;
- ii. treating physician will call LERN to request a transfer to another hospital;
- iii. LCC (LERN call center) will determine the closest and most appropriate facility available following LERN standard protocol;
- iv. if there are no available resources in the region then the LCC will locate an appropriate facility outside the region, and an interregional transfer will be considered. (All LERN interregional transfers will be reviewed by LERN medical directors and data will be collected for QI/PI.)

d. Exceptions

- i. EMS requesting LERN for patients located on or close to borders between two regions will and can be directed to either region based on the patient needs and available resources.
- ii. Air-med at the scene that is able to mitigate the time of transfer of long distances will and can be directed to hospitals outside the region they originate from based on patients needs and available resources.
- iii. LERN medical directors will be involved in the decision making (real time) in all patients that fall into the exception category.
- B. On August 20, 2009, the Louisiana Emergency Response Network Board [R.S. 40:2842(1)] adopted and promulgated the amended "Interregional Transfer Protocol" for the Louisiana Emergency Response Network [R.S. 40:2842(3)], as follows.
- 1. The LERN interregional transfer protocol only applies to those regions, hospitals and pre-hospital providers that are participating in the LERN network.

2. The interregional transfer protocol will be tested over a 90 day period, at the end of which all interregional transfers will be reviewed for compliance with protocols, quality, patient safety and standards of care. This information will be shared with regional commissions, LERN Board, and LERN design the system work group. Decisions regarding the Interregional Transfer Protocol will be made at the end of the 90-day trial period.

3. Interregional Transfer Protocol

- a. All patients whose conditions exceed the regionally available resources provided by local area hospitals may be transferred from one region to another following LERN interregional transfer protocol. Destination to the definitive care hospital in the receiving region will follow the LERN standard protocol. All laws regarding EMTALA apply.
- b. Only regions operating with the LERN standard protocol will be involved in the LERN interregional transfer protocol.
- c. Patients transferred via the LERN interregional transfer protocol must:
- i. be assessed at a local area hospital for treatment, be stabilized by a physician, and meet the entry criteria as determined by LERN standard protocol; and
- ii. have a treating physician call LERN to request a transfer to another hospital.
- d. The LERN call center (LCC) will determine the closest and most appropriate facility available following LERN standard protocol.
- e. If there are no available resources in the region, the LCC will locate an appropriate facility outside the region, and a interregional transfer will be considered.
- f. All LERN interregional transfers will be reviewed by LERN medical directors and data will be collected for QI/PI.

g. Exceptions

- i. Pre-hospital providers requesting LERN for patients located on or close to borders between regions will and can be directed to either region based on the patient needs and available resources.
- ii. Air-med at the scene able to mitigate the time of transfer of long distances will and can be directed to hospitals outside the region they originate from, based on patient needs and available resources.
- iii. LERN medical directors will be involved in the decision making for all patients in the exception category.
- C. These protocols were published at LR 35:2109-2110 (September 20, 2009).

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 41:141 (January 2015).

§19109. Standard LERN Entry and Destination Criteria

A. On October 21, 2010, the Louisiana Emergency Response Network Board [R.S. 40:2842(1)] revised, adopted and promulgated "Standard LERN Entry Criteria" and "Standard Destination Protocol" for use in all regions, except region 7, of the Louisiana Emergency Response Network [R.S. 40:2842(3)], replacing the "LERN Entry Criteria" and "LERN Region 4 LCC Destination Protocol," adopted and promulgated June 18, 2009, as follows.

1. Standard LERN Entry Criteria—Pre-Hospital and Hospital Triage Protocol

Standard LERN Entry Criteria			
Pre-Hospital and Hospital Triage Pro	tocol		
 □ Unmanageable Airway □ Tension Pneumothorax □ Traumatic cardiac arrest □ Burn patient without patent airway □ Burn patient >40 percent BSA without IV 	YES→	Call LCC	
Neurologic Trauma			
☐ GCS <14 + one or more of the following: ☐ Penetrating head injury or depressed skull fracture ☐ Open head injury with or without CSF leak ☐ Deterioration of the GCS ☐ Lateralizing signs or paralysis (i.e., one-sided weakness, motor, or sensory deficit) ■ Comparison of the following: ■ Comparison of the following:	YES→	Call LCC	
Physiologic			
□ SBP <90 (adults and > 9 y/o) <70 + 2 [age (yrs)] (age 1 to 8) <70 (age 1 to 12 months) <60 (term neonate) □ RR <10 or >29 (adults and ≥ 9 y/o) <15 or >30 (age 1 to 8) <25 or >50 (<12 m/o)	YES→	Call LCC	
Anatomic			
□ All penetrating injuries to neck, torso and extremities proximal to elbow and knee □ Flail Chest □ 2 or more proximal long-bone fractures □ Crush, degloved or mangled extremity □ Amputation proximal to wrist and ankle □ Pelvic Fracture □ Hip fractures (hip tenderness, deformity, lateral deviation of foot) excluding isolated hip fractures from same level falls □ Major joint dislocations (hip, knee, ankle, elbow) □ Open Fractures □ Fractures with neurovascular compromise (decreased peripheral pulses or prolonged capillary refill, motor or sensory deficits distal to fracture, etc.) Mechanism	YES→	Call LCC	
□ Falls > 20 ft. (adults)	YES→	Call LCC	
☐ Motorcycle crash >20 MPH			
Special ☐ Pregnancy >20 weeks ☐ Burns (will follow ABA guidelines)	YES→	Call LCC	

Standard LERN Entry Criteria			
Pre-Hospital and Hospital Triage Protocol			
Other			
\Box Age \geq 55 y/o or <8 y/o			
☐ Anticoagulation and bleeding disorders	YES→	Call	
☐ End stage renal disease	1 ES→	LCC	
☐ Transplant patients			

2. Standard Destination Protocol

□ Burn patient without patent airway □ Burn patient >40 percent BSA without IV NO ↓ Neurologic Trauma □ GCS <14 □ Penetrating head injury or depressed skull fracture □ Open head injury with or without CSF leak □ Deterioration of the GCS □ Lateralizing signs or paralysis (i.e., one-sided weakness, motor, or sensory deficit) NO ↓ Physiologic □ SBP <90 (adults and >9 y/o) <70 + 2 [age (yrs)] (age 1 to 8) <70 (age 1 to 12 months)	$ES \rightarrow$ $ES \rightarrow$	Closest ED		
□ Tension Pneumothorax □ Traumatic cardiac arrest □ Burn patient without patent airway □ Burn patient >40 percent BSA without IV NO Neurologic Trauma □ GCS <14 □ Penetrating head injury or depressed skull fracture □ Open head injury with or without CSF leak □ Deterioration of the GCS □ Lateralizing signs or paralysis (i.e., one-sided weakness, motor, or sensory deficit) NO Physiologic □ SBP <90 (adults and >9 y/o) <70 + 2 [age (yrs)] (age 1 to 8) <70 (age 1 to 12 months) <60 (term neonate) □ RR <10 or >29 (adults and ≥9 y/o) <15 or >30 (age 1 to 8) <25 or >50 (<12 m/o)	ES→	ED LERN Level I		
NO Neurologic Trauma GCS <14 Penetrating head injury or depressed skull fracture Open head injury with or without CSF leak Deterioration of the GCS Lateralizing signs or paralysis (i.e., one-sided weakness, motor, or sensory deficit) NO Physiologic SBP <90 (adults and >9 y/o) <70 + 2 [age (yrs)] (age 1 to 8) <70 (age 1 to 12 months) <60 (term neonate) RR <10 or >29 (adults and ≥9 y/o) <15 or >30 (age 1 to 8) <25 or >50 (<12 m/o)		Level I		
Neurologic Trauma □ GCS < 14 □ Penetrating head injury or depressed skull fracture □ Open head injury with or without CSF leak □ Deterioration of the GCS □ Lateralizing signs or paralysis (i.e., one-sided weakness, motor, or sensory deficit) NO ↓ Physiologic □ SBP < 90 (adults and > 9 y/o) < 70 + 2 [age (yrs)] (age 1 to 8) < 70 (age 1 to 12 months) < 60 (term neonate) □ RR < 10 or > 29 (adults and ≥ 9 y/o) < 15 or > 30 (age 1 to 8) < 25 or > 50 (< 12 m/o)		Level I		
GCS <14 □ Penetrating head injury or depressed skull fracture □ Open head injury with or without CSF leak □ Deterioration of the GCS □ Lateralizing signs or paralysis (i.e., one-sided weakness, motor, or sensory deficit) NO □ Physiologic □ SBP <90 (adults and >9 y/o) <70 + 2 [age (yrs)] (age 1 to 8) <70 (age 1 to 12 months) <60 (term neonate) □ RR <10 or >29 (adults and ≥9 y/o) <15 or >30 (age 1 to 8) <25 or >50 (<12 m/o)		Level I		
Penetrating head injury or depressed skull fracture □ Open head injury with or without CSF leak □ Deterioration of the GCS □ Lateralizing signs or paralysis (i.e., one-sided weakness, motor, or sensory deficit) NO ↓ Physiologic □ SBP <90 (adults and >9 y/o) <70 + 2 [age (yrs)] (age 1 to 8) <70 (age 1 to 12 months) <60 (term neonate) □ RR <10 or >29 (adults and ≥9 y/o) <15 or >30 (age 1 to 8) <25 or >50 (<12 m/o)		Level I		
NO ↓ Physiologic □ SBP <90 (adults and >9 y/o) <70 + 2 [age (yrs)] (age 1 to 8) <70 (age 1 to 12 months) <60 (term neonate)	ES→			
□ SBP <90 (adults and >9 y/o) <70 + 2 [age (yrs)] (age 1 to 8) <70 (age 1 to 12 months) <60 (term neonate) □ RR <10 or >29 (adults and ≥9 y/o) <15 or >30 (age 1 to 8) <25 or >50 (<12 m/o)	ES→			
□ SBP <90 (adults and >9 y/o) <70 + 2 [age (yrs)] (age 1 to 8) <70 (age 1 to 12 months) <60 (term neonate) □ RR <10 or >29 (adults and ≥9 y/o) <15 or >30 (age 1 to 8) <25 or >50 (<12 m/o)	ES→			
<pre></pre>	ES→			
NO		LERN Level I, II or III		
l I	NO			
↓				
Anatomic				
 All penetrating injuries to neck, torso and extremities proximal to elbow and knee Flail Chest 2 or more proximal long-bone fractures Crush, degloved or mangled extremity Amputation proximal to wrist and ankle Pelvic Fracture Hip fractures (hip tenderness, deformity, lateral deviation of foot) excluding isolated hip fractures from same level falls Major joint dislocations (hip, knee, ankle, elbow) Open Fractures Fractures with neurovascular compromise (decreased peripheral pulses or prolonged capillary refill, motor or sensory deficits distal to fracture, etc.) NO Mochanism				
Mechanism	Mechanism			
□ Falls > 20 ft. (adults)	ES→	LERN Level II or III		
or significant (>20 MPH) impact Motorcycle crash >20 MPH				

Standard Destination Protocol		
↓		
Special		
☐ Pregnancy >20 weeks		LERN
☐ Burns (will follow ABA guidelines)	$YES \rightarrow$	Level
		II or III
NO		
\downarrow		
Other		
\Box Age ≥ 55 y/o or <8 y/o		LERN
☐ Anticoagulation and bleeding disorders	MEG .	Level
☐ End stage renal disease	YES→	II, III
☐ Transplant patients		or IV

B. These protocols were published at LR 36:2743-2745 (November 20, 2010).

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 41:142 (January 2015).

§19111. Interregional Transfer Protocol

A. On January 20, 2011, the Louisiana Emergency Response Network Board (R.S. 40:2842(1) and (3)) adopted and promulgated "LERN Hospital Interregional Transfer Guidelines" and "LERN Hospital Interregional Transfer Protocol", replacing "Interregional Transfer Protocol" adopted June 18, 2009, as follows.

1. LERN Hospital Interregional Transfer Guidelines

- a. All patients whose conditions exceed the regionally available resources provided by local area hospitals may be transferred from one region to another following LERN interregional transfer protocol.
- b. The LERN hospital interregional transfer protocol only applies to hospitals that are participating in the LERN network.
- c. Regions or individual parishes that have MOU's (which include medical control and destination guidelines), between an ACS verified level 1 trauma center and a local parish medical society(ies) will be incorporated into the LCC standard operating procedure for the effected region(s).

2. LERN Hospital Interregional Transfer Protocol

- a. Patients transferred via the LERN hospital interregional transfer protocol must:
- i. meet LERN standard entry criteria that requires resources and/or capabilities not available in that region;
- ii. be assessed and stabilized to the best of their ability at a local area hospital prior to transport to the closest appropriate hospital;
- iii. the treating physician/nurse must contact LERN to request a transfer. The LERN communications center (LCC) will determine the closest and most appropriate facility available following the LERN standard destination protocol.

B. These guidelines and protocols were published at LR 37:751 (February 20, 2011).

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 41:143 (January 2015).

§19113. LERN Entry Criteria: Trauma; LERN Destination Protocol: Trauma

A. On January 20, 2011, the Louisiana Emergency Response Network Board [R.S. 40:2842(1) and (3)] adopted and promulgated "LERN ENTRY CRITERIA: Trauma; Pre-Hospital and Hospital Triage Protocol" and "LERN DESTINATION PROTOCOL: Trauma" replacing the "Standard LERN Entry Trauma Criteria" and "Standard LERN Entry Trauma Criteria Destination Protocol" adopted and promulgated January 20, 2011, as follows.

1. LERN Entry Criteria: Trauma

a. Pre-Hospital and Hospital Triage Protocol

	Call LERN Communications Center for:
	Unmanageable Airway
▮ •	Tension Pneumothorax
-	
•	Traumatic cardiac arrest
▮ •	Burn Patient without patent airway
Di	Burn patient >40 percent BSA without IV
Pn	ysiologic
•	GCS <14
•	SBP $<$ 90 (adults and $>$ 9 y/o)
	<70 (a = 1 to 12 m and a)
	<70 (age 1 to 12 months)
۱.	<60 (term neonate) RR <10 or >29 (adults & \geq 9 y/o)
•	<15 or >30 (age 1 to 8 y/o)
	<25 or >50 (<12 m/o)
Δn	natomic
•	Open or depressed skull fractures
	Open head injury with or without CSF leak
	Lateralizing signs or paralysis (i.e., one-sided weakness, motor, or
•	sensory deficit)
١.	All penetrating injuries to head, neck, torso, and extremities proximal
	to elbow and knee
	Flail Chest
	2 or more proximal long-bone fractures
	Crush, degloved or mangled extremity
	Amputation proximal to wrist and ankle
	Pelvic Fractures
	Hip Fractures (hip tenderness, deformity, lateral deviation of foot)
	excluding isolated hip fractures from same level falls
•	Major joint dislocations (hip, knee, ankle, elbow)
•	Open Fractures
•	Fractures with neurovascular compromise (decreased peripheral
	pulses or prolonged capillary refill, motor or sensory deficits distal to
	fracture)
Me	echanism
•	Falls >20 ft. adults
	>10 ft. (child) or 2 to 3 times height
•	Auto vs. pedestrian/bicyclist thrown, run over or significant
	(>20 MPH) impact
•	Motorcycle crash >20 MPH
•	High-risk auto crash
•	Intrusion >12 in, occupant site
	>18 in. any site
•	Ejection, partial or complete from automobile

Death in same passenger compartment

Call LERN Communications Center for: Other Pregnancy >20 weeks Burns (follow ABA guidelines) Age ≥ 55 y/o or <8 y/o Anticoagulation and bleeding disorders End stage renal disease Transplant patients

2. LERN Destination Protocol: Trauma

Multi/Mass Casualty Incident (MCI)

LERN Destination Protocol: Tra	uma	
Unmanageable Airway		
Tension Pneumothorax		Closest
Traumatic cardiac arrest	$YES \rightarrow$	ED
Burn patient without patent airway Burn patient > 40 paragent BSA without IV		
Burn patient >40 percent BSA without IV Physiologic		
• GCS < 14		
• SBP <90 (adults and > 9 y/o)		
<70 + 2 [age (yrs)] (age 1 to 8)		LEDN
<70 (age 1 to 12 months)	YES→	LERN Level I.
<60 (term neonate)	1E3→	II, or III
• RR <10 or >29 (adults and > 9 y/o)		11, 01 111
<15 or >30 (age 1 to 8)		
<25 or >50 (< 12 m/o)		
Anatomic Open or depressed skull fractures		
Open for depressed skull fractures Open head injury with or without CSF leak		
Lateralizing signs or paralysis (i.e., one-sided)		
weakness, motor, or sensory deficit)		
All penetrating injuries to neck, torso and		
extremities proximal to elbow and knee		
Flail Chest		
• 2 or more proximal long-bone fractures		
Crush, degloved or mangled extremity		LEDN
Amputation proximal to wrist and ankle	YES→	LERN Level I,
Pelvic Fracture Hip fractures (hip ton domness deformative)	1 E3→	II, or III
Hip fractures (hip tenderness, deformity, lateral deviation of foot) excluding isolated		11, 01 111
hip fractures from same level falls		
Major joint dislocations (hip, knee, ankle,		
elbow)		
Open Fractures		
Fractures with neurovascular compromise		
(decreased peripheral pulses or prolonged		
capillary refill, motor or sensory deficits distal to fracture		
Mechanism	l	l
□ Falls > 20 ft. (adults)		
> 10 ft. (child) or 2 to 3 times height		
☐ High-risk auto crash		
Intrusion > 12 in. occupant site:		LEDM
>18 in. any site	YES→	LERN Level II,
☐ Ejection, partial or complete from automobile	1 €5→	or III
☐ Death in same passenger compartment		() III
Auto vs. pedestrian/bicyclist thrown, run		
over or significant (>20 MPH) impact		
☐ Motorcycle crash >20 MPH		
Other Programmy > 20 weeks	1	
Pregnancy > 20 weeksBurns (follow ABA guidelines)		
Age >55 y/o or <8 y/o		LERN
 Age >33 y/0 of <8 y/0 Anticoagulation and bleeding disorders 	$YES \rightarrow$	Level II,
End stage renal disease		III, or IV
Transplant patients		
MULTI/MASS CASUALTY INCIDENT (MCI)		LERN
) ´	$YES \rightarrow$	Level I,
		II, III,

LERN Destination Protocol: Trauma	
	or IV

B. These protocols were published at LR 37:1466-1468 (April 20, 2011).

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 41:143 (January 2015).

§19115. LERN Destination Protocol: TRAUMA

A. On April 26, 2012, the Louisiana Emergency Response Network Board [R.S. 40:2842(1) and (3)] adopted and promulgated "LERN Destination Protocol: Trauma" replacing the "LERN Destination Protocol: Trauma" adopted and promulgated April 21, 2011, as follows.

Unmanageable Airway Tension Pneumothorax		Closest
Traumatic cardiac arrest	\rightarrow	ED
Burn Patient without patent airway		LD
Burn patient >40 percent BSA without IV		
Physiologic	l .	
• GCS <14		
• SBP <90 (adults and > 9 y/o)		
<70 + 2 [age (yrs)] (age 1 to 8 y/o)		
<70 (age 1 to 12 months)		Level I,
<60 (term neonate)	\rightarrow	II, or III*
• RR <10 or >29 (adults and ≥ 9 y/o)		
<15 or >30 (age 1 to 8 y/o)		
<25 or >50 (<12 m/o)		
Anatomic		
Open or depressed skull fractures		
Open head injury with or without CSF leak		
Lateralizing signs or paralysis (i.e., one-sided		
weakness, motor, or sensory deficit)		
All penetrating injuries to head, neck, torso, &		
extremities proximal to elbow & knee		
Flail Chest		
2 or more proximal long-bone fractures		
Crush, degloved or mangled extremity		
Amputation proximal to wrist & ankle	\rightarrow	Level I,
Pelvic Fractures	,	II, or III*
Hip Fractures (hip tenderness, deformity, lateral		
deviation of foot) excluding isolated hip fractures		
from same level falls		
 Major joint dislocations (hip, knee, ankle, 		
elbow)		
Open Fractures		
Fractures with neurovascular compromise		
(decreased peripheral pulses or prolonged capillary		
refill, motor or sensory deficits distal to fracture)		
Mechanism	1	
• Falls >20 ft. adults		
>10 ft. (child) or 2 to 3 times height		
High-risk auto crash		
 Intrusion > 12 in. occupant site 		
> 18 in. any site		Level I,
 Ejection, partial or complete from automobile 	\rightarrow	II, or III*
Death in same passenger compartment		
Auto vs. pedestrian/bicyclist thrown, run over or		
significant (>20 MPH) impact		
Motorcycle crash >20 MPH		
Other	I	
Other		

 Pregnancy >20 weeks Burns (follow ABA guidelines) Age ≥ 55 y/o or <8 y/o Anticoagulation and bleeding disorders -patients w/ head injuries are at high risk for rapid deterioration 	\rightarrow	Level II, or III*
MULTI/MASS CASUALTY INCIDENT (MCI)	\rightarrow	Level I, II, or III*

*Refers to ACS Verified Level Trauma Center—Where trauma center not available, patient will be routed to facility with appropriate resource which may not need be the highest level facility.

B. This protocol was published at LR 38:1462-1463 (June 20, 2012).

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 41:144 (January 2015).

§19117. LERN Destination Protocol: Trauma

A. On November 21, 2013, the Louisiana Emergency Response Network Board [R.S. 40:2842(1) and (3)] adopted and promulgated "LERN Destination Protocol: TRAUMA" replacing the "LERN Destination Protocol: Trauma" adopted and promulgated April 26, 2012, and repealing "LERN ENTRY CRITERIA, Trauma Pre-Hospital and Hospital Triage Protocol" adopted and promulgated April 21, 2011, as follows.

1. Call LERN Communication Center at (866) 320-8293 for patients meeting the following criteria.

Unmanageable airway Tension pneumothorax Traumatic cardiac arrest Burn patient without patent airway Burn patient > 40 percent BSA without IV	\rightarrow	Closest ED/Trauma Center
NO ↓		
Physiologic		
 GCS < 14 SBP < 90 (adults and > 9 y/o) < 70 + 2 [age (yrs)] (age 1 to 8 y/o) < 70 (age 1 to 12 months) < 60 (term neonate) RR < 10 or > 29 (adults & ≥ 9 y/o) < 15 or > 30 (age 1 to 8 y/o) < 25 or > 50 (< 12 m/o) 	\rightarrow	To Appropriate Trauma Center or Hospital as Determined by LERN Communication Center
NO ↓		

Anatomic		
 Open or depressed skull fractures Open head injury with or without CSF leak Lateralizing signs or paralysis (i.e., one-sided weakness, motor, or sensory deficit) All penetrating injuries to head, neck, torso, and extremities proximal to elbow and knee Flail Chest 2 or more proximal long-bone fractures Crush, degloved or mangled extremity Amputation proximal to wrist and ankle Pelvic Fractures Hip Fractures (hip tenderness, deformity, lateral deviation of foot) excluding isolated hip fractures from same level falls Major joint dislocations (hip, knee, ankle, elbow) Open Fractures Fractures with neurovascular compromise 	\rightarrow	To Appropriate Trauma Center or Hospital as Determined by LERN Communication Center
(decreased peripheral pulses or prolonged		
capillary refill, motor or sensory deficits distal to fracture)		
NO NO		
↓		
Mechanism		
Falls >20 ft. adults >10 ft. (child) or 2 to 3 times height High-risk auto crash Intrusion >12 in. occupant site >18 in. any site Ejection, partial or complete from automobile Death in same passenger compartment Auto vs. pedestrian/bicyclist thrown, run over or significant (>20 MPH) impact Motorcycle crash >20 MPH	\rightarrow	To Appropriate Trauma Center or Hospital as Determined by LERN Communication Center
NO ↓		
Other		
 Pregnancy >20 weeks Burns (follow ABA guidelines) Age ≥ 55 y/o or <8 y/o Anticoagulation and bleeding disorders - patients w/ head injuries are at high risk for rapid deterioration 	\rightarrow	To Appropriate Trauma Center or Hospital as Determined by LERN Communication Center
MULTI/MASS CASUALTY INCIDENT (MCI)	\rightarrow	To Appropriate Trauma Center or Hospital as Determined by LERN Communication Center

B. This protocol was published at LR 40:190-191 (January 20, 2014).

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Louisiana Emergency Response Network, LR 41:145 (January 2015).

§19119. Destination Protocol: TRAUMA

A. On November 20, 2014, the Louisiana Emergency Response Network Board [R.S. 40:2842(1) and (3)] adopted and promulgated "Destination Protocol: Trauma" to be

effective January 1, 2015, and replacing the "LERN Destination Protocol: Trauma" adopted and promulgated November 21, 2013, as follows.

1. Call LERN communication center at (866) 320-8293 for patients meeting the following criteria.

•	Unmanageable airway Tension pneumothorax Traumatic cardiac arrest Burn patient without patent airway Burn patient > 40 percent BSA without IV	Yes→	Closest ED/ Trauma Center
•	GCS ≤13 SBP <90mmHg RR <10 or >29 breaths per minute, or need for ventilator Support (<20 in infant aged <1 year)	Yes→	Transport to Trauma Center/ Trauma Program These patients should be transported to the highest level of care within the defined trauma system. This is a Level 1 or a Level 2 Trauma Center or Trauma Program. * If distance or patient condition impedes transport to trauma facility, consider transport to most appropriate resourced hospital.
	↓ No		
-	Assess anatomy of injury		m
•	All penetrating injuries to head, neck, torso, and extremities proximal to elbow or knee Chest wall instability or deformity (e.g. flail chest) Two or more proximal longbone fractures Crushed, degloved, mangled, or pulseless extremity Amputation proximal to wrist or ankle Pelvic fractures Open or depressed skull fracture Paralysis Fractures with neurovascular compromise (decreased peripheral pulses or prolonged capillary refill, motor or sensory deficits distal to fracture)	Yes→	Transport to Trauma Center/ Trauma Program These patients should be transported to the highest level of care within the defined trauma system. This is a Level 1 or a Level 2 Trauma Center or Trauma Program. * If distance or patient condition impedes transport to trauma facility, consider transport to most appropriate resourced hospital.
Α.	↓ No		
	sess mechanism of injury and idence of high-energy impact		
•	Falls - Adults: >20 feet (one story is equal to 10 feet) - Children: >10 feet or two or three times the height of the child High-risk auto crash - Intrusion, including roof: >12 inches occupant site;	Yes→	Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no Trauma Center/Trauma Program in the region, LCC may route to the

> 18 inches any site - Ejection (partial or complete) from automobile - Death in the same passenger compartment - Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/ATV thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph ↓ No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) most appropriate resourced hospital. most appropriate resourced hospital. resourced hospital. Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
complete) from automobile - Death in the same passenger compartment - Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/ ATV thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph ↓ No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms - Death in the same passenger compartment - Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
automobile - Death in the same passenger compartment - Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/ ATV thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph ↓ No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms
- Death in the same passenger compartment - Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/ ATV thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph ↓ No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms - Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/ ATV thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph ↓ No Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
passenger compartment - Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/ ATV thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph ↓ No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/ ATV thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph ↓ No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms - Vehicle telemetry data consistent with a high risk of injury Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/ ATV thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph ↓ No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms - Vehicle telemetry data consistent with a high risk of injury Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
consistent with a high risk of injury Auto vs. pedestrian/bicyclist/ ATV thrown, run over, or with significant (>20 mph) impact Motorcycle crash >20mph No Assess special patient or system considerations Older Adults Risk of injury/death increases after age 55 years SBP <110 may represent shock after age 65 Low impact mechanisms Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
risk of injury Auto vs. pedestrian/bicyclist/ ATV thrown, run over, or with significant (>20 mph) impact Motorcycle crash >20mph No Assess special patient or system considerations Older Adults Risk of injury/death increases after age 55 years SBP <110 may represent shock after age 65 Low impact mechanisms Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
Auto vs. pedestrian/bicyclist/ ATV thrown, run over, or with significant (>20 mph) impact Motorcycle crash >20mph
ATV thrown, run over, or with significant (>20 mph) impact Motorcycle crash >20mph No Assess special patient or system considerations Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
significant (>20 mph) impact Motorcycle crash >20mph No Assess special patient or system considerations Older Adults Risk of injury/death increases after age 55 years SBP <110 may represent shock after age 65 Low impact mechanisms Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
Motorcycle crash >20mph
↓ No Assess special patient or system considerations Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms Assess special patient or system Crenter/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
Assess special patient or system considerations Older Adults Risk of injury/death increases after age 55 years SBP <110 may represent shock after age 65 Low impact mechanisms Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
considerations Older Adults Risk of injury/death increases after age 55 years SBP <110 may represent shock after age 65 Low impact mechanisms Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
Older Adults Risk of injury/death increases after age 55 years SBP <110 may represent shock after age 65 Low impact mechanisms Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
- Risk of injury/death increases after age 55 years which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no
- SBP <110 may represent shock after age 65 system, need not be the highest level trauma center/program. If no
- SBP <110 may represent shock after age 65 shock ammachanisms system, need not be the highest level trauma center/program. If no
shock after age 65 highest level trauma - Low impact mechanisms center/program. If no
- Low impact mechanisms center/program. If no
i te o oronna jever rans) i i ranna Cenier/ Franna
may result in severe Program in the region,
injury LCC may route to the
Should be thaged
preferentially to pediatric
capable trauma centers
Anticoagulants and bleeding
disorders Yes→
- Patients with head injury
are at high risk for rapid
deterioration
Burns
- With trauma mechanism:
triage to trauma center
Pregnancy >20 weeks
Hip Fractures (hip tenderness,
deformity, lateral deviation of
foot) excluding isolated hip
fractures from same level falls
Major joint dislocations (hip,
, 1.
knee, ankle, elbow)
Open Fractures
EMS provider judgment
↓No
Multi/Mass Casualty Incident No→ Transport according to
protocol

2. When in doubt, transport to a trauma center.

B. This protocol was published at LR 40:2710 (December 20, 2014).

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 41:950 (May 2015).

§19121. LERN Destination Protocol: TRAUMA

A. On December 10, 2015, the Louisiana Emergency Response Network Board [R.S. 40:2842(1) and (3)] adopted and promulgated "LERN Destination Protocol: TRAUMA", which replaces the "LERN Destination Protocol: TRAUMA" found in §19121 adopted and promulgated November 20, 2014, as follows.

1. Call LERN communication center at (866) 320-8293 for patients meeting the following criteria.

8293 for patients meeting t	ne ron	owing criteria.
Unmanageable airway Tension pneumothorax Traumatic cardiac arrest Burn patient without patent airway Burn patient > 40 percent BSA without IV	Yes→	Closest ED/Trauma Center
No		
Measure vital signs and		
level of consciousness		
GCS ≤13 SBP <90mmHg RR <10 or >29 breaths per minute, or need for ventilator Support (<20 in infant aged <1 year)	Yes→	Transport to Trauma Center/ Trauma Program These patients should be transported to the highest level of care within the defined trauma system. This is a Level 1 or a Level 2 Trauma Center or Trauma Program. * If distance or patient condition impedes transport to trauma facility, consider transport to most appropriate resourced hospital.
No		
Aggagg anotamy of::		
• All penetrating injuries to		
An penetrating injuries to head, neck, torso, and extremities proximal to elbow or knee Chest wall instability or deformity (e.g. flail chest) Two or more proximal long-bone fractures Crushed, degloved, mangled, or pulseless extremity Amputation proximal to wrist or ankle Pelvic fractures Open or depressed skull fracture Paralysis Fractures with neurovascular compromise (decreased peripheral pulses or prolonged capillary refill, motor or sensory deficits distal to fracture)	Yes→	Transport to Trauma Center/ Trauma Program These patients should be transported to the highest level of care within the defined trauma system. This is a Level 1 or a Level 2 Trauma Center or Trauma Program. * If distance or patient condition impedes transport to trauma facility, consider transport to most appropriate resourced hospital.
↓ ↓		
Assess mechanism of injury and evidence of high-energy impact		
Falls Adults: >20 feet (one story is equal to 10 feet) Children: >10 feet or two or three times the height of the child High-risk auto crash Intrusion, including roof: > 12 inches occupant site; > 18 inches any site Ejection (partial or complete) from automobile	Yes→	Transport to Trauma Center/Trauma Program which, depending upon the defined trauma system, need not be the highest level trauma center/program. If no Trauma Center/Trauma Program in the region, LCC may route to the most appropriate resourced hospital.

passenger compartment - Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury • Children - Should be triaged preferentially to pediatric capable trauma centers - Patients with head injury are at high risk for rapid deterioration • Burns - With trauma mechanism: triage to trauma center • Pregnancy >20 weeks • Hip Fractures (hip tenderness, deformity, lateral deviation of foot) excluding isolated hip fractures from same level falls • Major joint dislocations (hip, knee, ankle, elbow) • Open Fractures • EMS provider judgment No	Multi/Mass Casualty Incident	No→	Transport according to protocol
- Vehicle telemetry data consistent with a high risk of injury - Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact - Motorcycle crash >20mph No Assess special patient or system considerations - Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury - Children - Should be triaged preferentially to pediatric capable trauma centers - Anticoagulants and bleeding disorders - Patients with head injury are at high risk for rapid deterioration - Burns - With trauma mechanism: triage to trauma center - Pregnancy >20 weeks - Hip Fractures (hip tenderness, deformity, lateral deviation of foot) excluding isolated hip fractures from same level falls - Major joint dislocations (hip, knee, ankle, elbow) - Open Fractures - EMS provider judgment	140		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury • Children - Should be triaged preferentially to pediatric capable trauma centers • Anticoagulants and bleeding disorders - Patients with head injury are at high risk for rapid deterioration • Burns - With trauma mechanism: triage to trauma center • Pregnancy >20 weeks • Hip Fractures (hip tenderness, deformity, lateral deviation of foot) excluding isolated hip fractures from same level	 Major joint dislocations (hip, knee, ankle, elbow) Open Fractures 		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury • Children - Should be triaged preferentially to pediatric capable trauma centers • Anticoagulants and bleeding disorders - Patients with head injury are at high risk for rapid deterioration • Burns - With trauma mechanism: triage to trauma center • Pregnancy >20 weeks	tenderness, deformity, lateral deviation of foot) excluding isolated hip fractures from same level		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury • Children - Should be triaged preferentially to pediatric capable trauma centers • Anticoagulants and bleeding disorders - Patients with head injury are at high risk for rapid deterioration • Burns - With trauma mechanism: triage to trauma center			
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No ↓ Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury • Children - Should be triaged preferentially to pediatric capable trauma centers • Anticoagulants and bleeding disorders - Patients with head injury are at high risk for rapid deterioration • Burns - With trauma			
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No ↓ Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury • Children - Should be triaged preferentially to pediatric capable trauma centers • Anticoagulants and bleeding disorders - Patients with head injury are at high risk for rapid deterioration • Burns Transport to Trauma Center/Trauma Program or hospital capable of timely and thorough evaluation and initial management of potentially serious injuries. Consider consultation with medical			
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury • Children - Should be triaged preferentially to pediatric capable trauma centers • Anticoagulants and bleeding disorders - Patients with head injury are at high risk for rapid deterioration Transport to Trauma Center/Trauma Program or hospital capable of timely and thorough evaluation and initial management of potentially serious injuries. Consider consultation with medical			control.
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury • Children - Should be triaged preferentially to pediatric capable trauma centers • Anticoagulants and bleeding disorders - Patients with head injury are at high risk for rapid Transport to Trauma Center/Trauma Program or hospital capable of timely and thorough evaluation and initial management of potentially management of potentially			
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury • Children - Should be triaged preferentially to pediatric capable trauma centers • Anticoagulants and bleeding disorders - Patients with head injury Ves→ Transport to Trauma Center/Trauma Program or hospital capable of timely and thorough evaluation and initial	are at high risk for rapid		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury • Children - Should be triaged preferentially to pediatric capable trauma centers • Anticoagulants and		Yes→	thorough evaluation and initial
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury • Children - Should be triaged preferentially to pediatric capable trauma centers Transport to Trauma Contor/Trauma Program or	S		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury • Children - Should be triaged	pediatric capable trauma centers		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may result in severe injury	- Should be triaged		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g. ground level falls) may	3 2		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65 - Low impact mechanisms (e.g.			
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent shock after age 65	mechanisms (e.g.		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55 years - SBP <110 may represent			
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death increases after age 55			
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults - Risk of injury/death	_		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations • Older Adults			
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No Assess special patient or system considerations			
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph No No ↓	system considerations		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash >20mph	Assess special patient or		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact • Motorcycle crash	No		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 mph) impact	·		
 Vehicle telemetry data consistent with a high risk of injury Auto vs. pedestrian/bicyclist/AT V thrown, run over, or with significant (>20 			
 Vehicle telemetry data consistent with a high risk of injury Auto vs. pedestrian/bicyclist/AT V thrown, run over, or 	· ·		
- Vehicle telemetry data consistent with a high risk of injury • Auto vs.			
- Vehicle telemetry data consistent with a high risk of injury			
- Vehicle telemetry data consistent with a high	, , , , , , , , , , , , , , , , , , ,		
- Vehicle telemetry data			
passenger compartment			
- Death in the same			

- 2. When in doubt, transport to a trauma center.
- B. This protocol was published at LR 42:169 (January 2016).

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 42:904 (June 2016).


Chapter 192. Burn Protocols

§19201. LERN Destination Protocol: BURN

A. Call LERN Communication Center at 1-866-320-8293 for patients meeting the following criteria.

Burn Patient with Trauma	Yes→	See LERN Trauma Destination Protocol
↓No		
Burn patient without patent airway Patients with facial / airway	Yes→	Transport to Closest ED
burns or anticipated airway compromise Burn patient with > 40 % BSA without IV or IO access		
↓No		
2 nd and 3 rd degree burns involving: > 10% BSA Face, hands, feet, genitalia, perineum, or major joints Circumferential Burns Electrical burns, including lightning injury Chemical burns Or Inhalation injury All Third Degree Burns	Yes→	Transport to Closest Burn Center * If distance or patient condition impedes transport to burn center, consider transport to most appropriate resourced hospital.
Transport per Local EMS Protocols		

- In the event of a burn disaster, each burn center should immediately contact LERN. LERN Call Center (LCC) will conduct a bed poll.
- In the event of a burn disaster and excess beyond capacity, the next geographically closest burn center should be alerted immediately by LERN.



AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2846(A), R.S. 40:2845(A)(1) and R.S. 9:2798.5.

HISTORICAL NOTE: Promulgated by the Department of Health, Emergency Response Network, LR 45:911 (July 2019).

Chapter 193. Stroke Protocols

§19301. LERN Destination Protocol: Stroke

- A. On November 21, 2013, the Louisiana Emergency Response Network Board [R.S. 40:2842(1) and (3)] adopted and promulgated "LERN Destination Protocol: STROKE," as follows.
- 1. The following protocol applies to patients with suspected stroke.

→ Stroke Level I, II,	•		
All other patients with suspected stroke Patients with seizure with focal deficit, extended window (4-8 hrs from onset), and patients with unknown onset may benefit from evaluation at Level I or II hospital with on-site stroke expertise. NO ↓ Terminally Ill or Palliative Care Patient Transport to LERN Stroke Level I, II, or III Transport to LERN Stroke Level I, II,	AirwayBreathingCirculation	\rightarrow	Closest ED
Patients with seizure with focal deficit, extended window (4-8 hrs from onset), and patients with unknown onset may benefit from evaluation at Level I or II hospital with on-site stroke expertise. NO Terminally Ill or Palliative Care Patient Patients with focal deficit prometics with unknown onset may benefit from evaluation at Level I or II hospital Level I, II, or III Transport to LERN Stroke Level I, II, or III	NO ↓		
	Patients with seizure with focal deficit, extended window (4-8 hrs from onset), and patients with unknown onset may benefit from evaluation at Level I or II hospital	\rightarrow	LERN Stroke
→ Stroke Level I, II,	NO ↓		
Cuiding principles:	,	\rightarrow	Transport to LERN Stroke Level I, II, III, or IV

Guiding principles:

- · Time is the critical variable in acute stroke care.
- Protocols that include pre-hospital notification while en route by EMS should be used for patients with suspected acute stroke to facilitate primary destination efficiency.
- Treatment with intravenous tPA is the only FDA approved acute therapy for stroke.
- EMS should identify the geographically closest facility capable of providing tPA treatment.
- Transfer patient to the nearest hospital equipped to provide tPA treatment.
- Secondary transfer to facilities equipped to provide tertiary care and interventional treatments should not prevent administration of tPA to appropriate patients.
- B. This protocol was published at LR 40:189-190 (January 20, 2014).

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).


HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 41:146 (January 2015).

§19303. LERN Destination Protocol: Stroke

A. On April 21, 2017, the Louisiana Emergency Response Network Board [R.S. 40:2842(1) and (3)] adopted and promulgated "LERN Designation Protocol: Stroke", amending and replacing the previous "LERN Designation Protocol: Stroke" adopted on November 21, 2013 and set out in Section 19301, as follows.

LERN Destination Protocol: Stroke LERN Call Center: (866) 320-8293

The following protocol applies to patients with suspected stroke:



All other patients with suspected stroke, determine time last seen normal (LSN) and screen for large vessel occlusion (LVO)



LSN < 6 hours* AND screen for LVO is positive

Transport to LERN Stroke Level I, II, or III Center

If < 15 minutes of additional transport time to reach Level I or endovascular capable Level II Center, transfer to the Level I or endovascular capable Level II Center

LSN > 6 hours OR screen for LVO is negative

Transport to LERN Stroke Level I, II, or III Center

If > 15 minutes of additional transport time to reach Level I, II, or III Center than to reach stroke capable Off Site ED, it is acceptable to transport to a stroke capable Off Site

* The LSN < 6hrs should include patients without a definite time of LSN, but who could reasonably be assumed to be within 6 hrs of onset, including patients who wake-up with stroke symptoms.

Guiding Principles:

- Time is the critical variable in acute stroke care
- Protocols that include pre-hospital notification while en route by EMS should be used for patients with suspected acute stroke to facilitate initial destination efficiency.
- Treatment with intravenous tPA is the only FDA approved medication therapy for hyperacutestroke.
- EMS should identify the geographically closest hospital capable of providing tPA treatment.
- Transfer patient to the nearest hospital equipped to provide tPA treatment.
- Secondary transfer to facilities equipped to provide tertiary care and interventional treatments should not prevent administration of tPA to appropriate patients.

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health, Emergency Response Network, LR 43:1758 (September 2017).

Chapter 195. STEMI Protocols

§19501. STEMI Triage Protocol for Pre-Hospital Providers

A. On November 21, 2013, the Louisiana Emergency Response Network Board [R.S. 40:2842(1) and (3)] adopted and promulgated "STEMI Triage Protocol for Pre-Hospital Providers," as follows.

Acute coronary symptoms ≥ 15 minutes and < 12 hours

AND

12 lead ECG criteria of 1 mm ST elevation in 2 or more contiguous leads

OR

LBBB NOT KNOWN to be present in the past

EMS ECG interpreted or transmitted to hospital for MD consult for bypass and activation

↓

STEMI-Receiving Center with medical contact-to-device (PCI) ≤ 90 minutes (by ground or air)?	YES→	Transport to nearest STEMI-Receiving Center with pre-hospital notification/activation Goal medical contact to device (PCI) time of 90 minutes or less
NO ↓		
Transport to closest STEMI-Referral Hospital with Pre- hospital notification/activation Goal medical contact to fibrinolytic needle time of 30 minutes or less	\rightarrow	Transport to nearest STEMI-Receiving Center for subsequent PCI

*O'Gara PT, Kushner FG, Ascheim DD, et all. 2013 ACCF/AHA Guideline for the Management of ST-Elevation Myocardial Infraction: A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. Journal of the American College of Cardiology. 2013;61(4):e78.

B. This protocol was published at LR 50:192 (January 20, 2014).

AUTHORITY NOTE: Promulgated in accordance with R.S. 9:2798.5 and R.S. 40:2846(A).

HISTORICAL NOTE: Promulgated by the Department of Health and Hospitals, Emergency Response Network, LR 41:146 (January 2015).

Chapter 197. Trauma Program Recognition

§19701. Generally

A. The goal of the Louisiana Emergency Response Network Board is to establish a trauma system that includes one verified trauma center in each region of the state. Trauma program recognition in excess of this goal will be determined utilizing a needs based assessment. The LERN communication center coordinates access to the trauma system by providing accurate and professional routing of patients experiencing time sensitive illness to the definitive care facility, which includes trauma programs recognized according to these rules.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2846(A), R.S. 40:2845(A)(1) and R.S. 9:2798.5.

HISTORICAL NOTE: Promulgated by the Department of Health, Emergency Response Network, LR 42:1931 (November 2016).

§19703. Purpose

A. LERN recognizes the opportunity to reduce the morbidity and mortality of trauma patients in Louisiana in areas without an existing level I or level II trauma center or an existing level II or level III trauma program through this process which recognizes the achievement of specific benchmarks in hospitals actively pursuing levels II or III trauma center verification through the American College of Surgeons (ACS).

- B. The purpose of this Chapter is to define the qualifications, procedure, and requirements for hospitals seeking trauma center verification by the ACS to be recognized by LERN as achieving the core components of a trauma program and thus qualified for recognition as a trauma program.
- C. The criteria for trauma program recognition are drawn from *Resources for Optimal Care of Injured Patient 2014* published by the ACS.
- D. Trauma program recognition is distinct and different from the trauma center certification by the state. To be certified as a trauma center, a hospital must satisfy the requirements of R.S. 40:2172 and 2173.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2846(A), R.S. 40:2845(A)(1) and R.S. 9:2798.5.

HISTORICAL NOTE: Promulgated by the Department of Health, Emergency Response Network, LR 42:1931 (November 2016).

§19705. Qualifications for LERN Trauma Program Recognition

- A. The hospital must be located in a LERN region that does not have an existing ACS verified level I or level II trauma center.
- B. A hospital providing care to trauma patients in a LERN region without an existing ACS verified level I or level II trauma center or without an existing level II or level III trauma program is eligible for trauma program recognition upon meeting the requirements of this rule.
- C. If there is an existing LERN recognized level II or Level III trauma program in the LERN region, the hospital must complete the most current version of the ACS needs based assessment of trauma systems tool (ACS NBATS). If the number of trauma centers allocated by the tool is less than or equal to the number of existing trauma programs in the region, the hospital is not eligible for trauma program recognition.
- D. A hospital must be in the process of working toward ACS verification to be eligible for trauma program recognition.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2846(A), R.S. 40:2845(A)(1) and R.S. 9:2798.5.

HISTORICAL NOTE: Promulgated by the Department of Health, Emergency Response Network, LR 42:1932 (November 2016).

§19707. Procedure for Trauma Program Recognition

- A. A hospital must complete the LERN approved form, "application for recognition of trauma program".
- B. The hospital CEO must complete and sign the LERN approved trauma program checklist/attestation for the applicable trauma program level.
- 1. By this attestation, the hospital CEO ensures 24/7/365 availability of the resources listed.
- 2. The attestation must be validated by a site visit by LERN staff.

- 3. Upon CEO attestation and/or site visit, if it is determined by the LERN executive committee in conjunction with the LERN trauma medical director, that the required benchmarks are not in place the hospital will not be eligible for trauma program verification.
- C. After satisfying the requirements of A. and B. above, the hospital will be recognized as a trauma program and such recognition will be added to the LERN resource management screen for the purpose of routing trauma patients.
- D. To maintain trauma program recognition, the hospital must request an ACS verification or consultation site visit at the time of the attestation or within 30 days thereafter, with the consultation or survey to occur within 12 months of the attestation or as close to 12 months as the ACS schedule allows. Written documentation of the request and scheduling must be submitted to LERN.
- 1. If an ACS verification or consultation site visit is not requested within 30 days and does not occur within 12 months or as close to 12 months as the ACS schedule allows, the trauma program indicator on LERN resource management screen will be removed.
- E. After a consultation visit for the desired trauma level, the hospital has 30 days to schedule the verification survey by the ACS to occur within 12 months of the consultation or as close to 12 months as the ACS schedule allows. Written documentation of the request and scheduling must be submitted to LERN.
- 1. If documentation of scheduling per required parameters is not submitted to LERN and the ACS verification survey is not scheduled to occur within 12

- months of the consultation or as close to 12 months as the ACS schedule allows, the trauma program indicator will be removed on the LERN resource management screen.
- 2. If the hospital fails the ACS verification visit and a focused review visit, the hospital will lose trauma program status. The trauma program indicator will be removed on the LERN resource management screen.
- F. After loss of trauma program status for failing the ACS verification visit and focused review visit, trauma program status may be regained provided the following conditions are met:
- 1. a LERN designee and either the LERN trauma medical director or a trauma surgeon must review the deficiencies and findings of the ACS at a site visit;
- 2. the hospital must develop a remediation plan and apply to the LERN board for approval of trauma program status;
- 3. the LERN board will review the LERN team assessment of deficiencies and the hospital's remediation plan;
- 4. the LERN board must vote to approve the trauma program status request.

AUTHORITY NOTE: Promulgated in accordance with R.S. 40:2846(A), R.S. 40:2845(A)(1) and R.S. 9:2798.5.

HISTORICAL NOTE: Promulgated by the Department of Health, Emergency Response Network, LR 42:1932 (November 2016), amended LR 44:63 (January 2018), LR 45:436 (March 2019), repromulgated LR 45:573 (April 2019).